

The evolution of atomic matter waves inside an atom fiber

Jian Qi Shen^{1,2 *}

¹ Centre for Optical and Electromagnetic Research, Joint Research Centre of Photonics of The Royal Institute of Technology (Sweden) and Zhejiang University, Zhejiang University, Hangzhou Yuquan 310027, P.R. China

² Zhejiang Institute of Modern Physics and Department of Physics, Zhejiang University, Hangzhou 310027, P.R. China
(February 1, 2008)

A phenomenological description of time evolution of atomic matter waves inside a spiral shaped atomic-wave guide is presented in this report. We study three related topics: (i) the effective Hamiltonian and the time-development equation governing the matter waves; (ii) wavefunctions describing the matter waves in the noncoplanar atom guide; (iii) showing that such wavefunctions obtained is just the eigenstates of the atomic momentum operator. It is believed that both the idea and the results presented here may have relevance to some fundamental problems of *atom optics*.

PACS number(s): 03.65.Ca, 32.80.Pj

Recently, the implementation of atom chips based on the microfabrication techniques [1–3] in atom optics can allow trapping and guiding of atoms with a high accuracy, and therefore has many applications [4–6] in mesoscopic physics. More recently, Luo *et al.* reported a novel work of the so-called “atom fiber”, *i.e.*, an atomic guide that can control the matter waves (such as cold neutral atoms and ideally Bose-Einstein condensates) on a two dimensional surface of the atom chip [7]. In their experiments, Luo *et al.* demonstrated guiding atoms around more than two complete turns along a spiral shaped 25 mm long curved path at various atom-surface distances [7]. Such a technique is a key element for the controlled manipulation of matter waves on the atom chip. It is believed that the atom fiber and related technologies may have some potential applications to, *e.g.*, the quantum registers for quantum information processing and the guided matter wave interferometers (*e.g.*, the wide angle Mach-Zehnder and Sagnac interferometers) [7].

In Luo *et al.*’s paper, investigators presented the experimental details and mechanism for the realization of the atomic matter-wave guide [7]. Because of the complexity of the conditions for realizing the atom fiber, the theoretical description of wave propagation of atoms inside it can be achieved neither in a straightforward way nor from the first principle. In this report we demonstrate that based on some assumptions, which can simplify the interaction details of atomic matter wave on the atom chip, there may exist a phenomenological description that will give an enlightening understanding of the time evolution of matter waves inside the curved atomic wave guide.

In what follows, by ignoring the subordinate details of state evolution in an atom fiber, we assume that the atom fiber can be viewed as a perfect guide, namely, (i) the guide does not alter the magnitude but the direction of atom wave vector $\mathbf{k}$ in the evolution process, and (ii) from the point of view of mesoscopic scale, the wave vector $\mathbf{k}$ of atomic matter waves may be said to be always along the tangent to the curved guide (path) at each point at arbitrary time in the evolution process inside the atom fiber. Note that here the second assumption can truly hold in this process since, for example, in the side guide that is the simplest form of a magnetic guide, atoms can only move along the third unconfined direction as long as the external bias field points in the direction perpendicular to the guiding direction [7]. In this sense, the wave vector of atoms can characterize the geometric shape of the wire guide, which enables us to deal with the propagation problem of atomic matter waves in the spiral shaped wire guide. The aim of this report is to develop a phenomenological method for treating the time evolution of atom wavefunctions based on these two assumptions in such atom fibers.

According to the above two assumptions, the momentum (wave vector) squared $\mathbf{k}^2$ of a matter wave propagating inside a noncoplanarly curved atom fiber is conserved, which will yield the following identity

$$\dot{\mathbf{k}} + \mathbf{k} \times \left(\frac{\mathbf{k} \times \dot{\mathbf{k}}}{k^2} \right) = 0, \quad (1)$$

where dot denotes the derivative with respect to time, and k is the magnitude of $\mathbf{k}$, *i.e.*, $k = |\mathbf{k}|$. Although Luo *et al.*’s atom fiber was the coplanar shaped wire guide on the chip surface (*i.e.*, the spiral guild has a two-dimensional wire plane), instead, here we will consider a more general case in which the atom fiber is noncoplanarly shaped. This,

*E-mail address: jqshen@coer.zju.edu.cn

therefore, means that we will not restrict the atom wave vector only to the two-dimensional spiral shaped fiber (*i.e.*, in the following discussion $\mathbf{k}$ can be a three dimensional vector). Since Eq.(1) is exactly analogous to the equation of motion of a charged particle moving in a magnetic field or of a spinning particle in a rotating frame of reference, $\mathbf{k} \times \dot{\mathbf{k}}/k^2$ can be regarded as a “magnetic field” or “gravitomagnetic field” [8,9] (correspondingly, $\mathbf{k} \times (\mathbf{k} \times \dot{\mathbf{k}})/k^2$ can be thought of as a “Lorentz magnetic force” or “Coriolis force”). Just similar to Mashhoon’s work of deriving the interaction Hamiltonian of gravitomagnetic dipole moment (*i.e.*, spin) in a gravitomagnetic field [10,11], one can also readily obtain the Hamiltonian describing the coupling of the atomic total “gravitomagnetic moment” (*i.e.*, the total angular momentum operator $\mathbf{J}$) to the “gravitomagnetic field” as follows

$$H(t) = \frac{\mathbf{k}(t) \times \dot{\mathbf{k}}(t)}{k^2} \cdot \mathbf{J}, \quad (2)$$

which may be the effective Hamiltonian describing the propagation of atomic matter waves in the atom fiber.

The above derivation of the effective Hamiltonian is enlightening, the physical meanings of which is explicit. Now we will treat this problem in a more rigorous way, *i.e.*, by using the infinitesimal rotation operator of atom wavefunction. The atom wavefunction $|m, \mathbf{k}(t)\rangle$ (with m being the eigenvalue of the third component of $\mathbf{J}$) varies as it rotates by an infinitesimal angle, say $\vec{\vartheta}$, namely, it obeys the following transformation rule (in the unit $\hbar = 1$)

$$|m, \mathbf{k}'(t)\rangle = \exp \left[-i\vec{\vartheta} \cdot \mathbf{J} \right] |m, \mathbf{k}(t)\rangle, \quad (3)$$

where $\exp \left[-i\vec{\vartheta} \cdot \mathbf{J} \right] \simeq 1 - i\vec{\vartheta} \cdot \mathbf{J}$, and $\mathbf{k}'(t) = \mathbf{k}(t) + \Delta\mathbf{k}(t)$ with $\Delta\mathbf{k}(t) = \dot{\mathbf{k}}\Delta t$. Here $|\vec{\vartheta}|$ is the angle between $\mathbf{k}(t)$ and $\mathbf{k}'(t)$, and the direction of $\vec{\vartheta}$ is parallel to that of $\mathbf{k}(t) \times \mathbf{k}'(t)$. One can therefore arrive at

$$\vec{\vartheta} = \frac{\mathbf{k}(t) \times \mathbf{k}'(t)}{k^2} = \frac{\mathbf{k}(t) \times \dot{\mathbf{k}}(t)}{k^2} \Delta t. \quad (4)$$

Expanding $|m, \mathbf{k}'(t)\rangle$ in (3) to the first order, it follows that

$$i \frac{\partial |m, \mathbf{k}(t)\rangle}{\partial t} = \frac{\mathbf{k}(t) \times \dot{\mathbf{k}}(t)}{k^2} \cdot \mathbf{J} |m, \mathbf{k}(t)\rangle. \quad (5)$$

Not that Eq.(5) is just the form of the time-dependent Schrödinger equation, which governs the evolution of atomic matter waves inside a noncoplanar atom fiber. Thus by using the technique of infinitesimal rotation of atom wavefunctions, we obtain the effective Hamiltonian (2) (and hence the time-dependent Schrödinger equation) of atoms moving inside the curved atomic-wave guild.

It follows from Eq.(5) that the atom fiber with atomic matter waves inside it is just the three-generator Hamiltonian system, the interaction Hamiltonian of which is the linear combination of three algebraic generators which form a certain Lie algebra, say, $su(2)$ for the present system. In the literature, we have obtained the exact solutions of the arbitrary three-generator time-dependent quantum systems (and models) [12]. Based on this work [12], here we will straightforwardly write the wavefunctions of Eq.(5) without presenting the details of derivation procedure. For convenience, in what follows the atom wave vector is rewritten in the spherical polar coordinate system, *i.e.*, $\mathbf{k}(t) = k(\sin \theta \cos \varphi, \sin \theta \sin \varphi, \cos \theta)$, where both θ and φ are the time-dependent functions. We assume that the initial wave vector is $\mathbf{k}(0) = (0, 0, k)$, *i.e.*, the initial polar angle $\theta(0) = 0$. According to the results obtained in the reference [12], the wavefunction in Eq.(5) can be written in the form

$$|m, \mathbf{k}(t)\rangle = \exp \left[\frac{1}{i} \phi_m(t) \right] V(t) |m\rangle, \quad (6)$$

where $|m\rangle$ is the initial state satisfying the eigenvalue equation $J_3|m\rangle = m|m\rangle$, and the expression for the phase is $\phi_m(t) = m \int_0^t \dot{\varphi}(t') [1 - \cos \theta(t')] dt'$. The time-dependent operator $V(t)$ takes the form

$$V(t) = \exp [\beta(t)J_+ - \beta^*(t)J_-] \quad (7)$$

with $\beta = -(\theta/2) \exp(-i\varphi)$ and $\beta^* = -(\theta/2) \exp(i\varphi)$. Here the operators $J_{\pm} = J_1 \pm iJ_2$.

It is apparently seen that the momentum operator $\mathbf{p}$ of atoms is also a physical quantity that can characterize the propagation of atomic matter waves in the coiled atom fiber, since in accordance with the second assumption, the direction of wave vector $\mathbf{k}(t)$ (the time-dependent eigenvalue of $\mathbf{p}$) contains the information on the geometric shape

of the curved wire guide. In order to show that our treatment of evolution of atomic matter waves in the atom fiber is self-consistent, we should prove that the above wavefunction is the eigenstate of the momentum operator of atoms in the evolution process. In the Cartesian coordinate system, by the aid of the following commuting relations

$$[J_i, p_j] = i\epsilon_{ijk}p_k \quad (8)$$

with ϵ_{ijk} being the three dimensional Levi-Civita tensor, one can arrive at

$$\begin{aligned} V^\dagger(t)p_1V(t) &= p_1 + \sin\theta \cos\varphi p_3 + \frac{(1 - \cos\theta) \cos\varphi}{\theta} (\beta p_+ + \beta^* p_-), \\ V^\dagger(t)p_2V(t) &= p_2 + \sin\theta \sin\varphi p_3 + \frac{(1 - \cos\theta) \sin\varphi}{\theta} (\beta p_+ + \beta^* p_-), \\ V^\dagger(t)p_3V(t) &= p_3 + (\cos\theta - 1)p_3 + \frac{\sin\theta}{\theta} (\beta p_+ + \beta^* p_-), \end{aligned} \quad (9)$$

where $p_\pm = p_1 \pm ip_2$. With the help of these relations, one can obtain

$$p_i|m, \mathbf{k}(t)\rangle = [k \sin\theta(\delta_{1i} \cos\varphi + \delta_{2i} \sin\varphi) - k(1 - \cos\theta)\delta_{3i} + k_i]|m, \mathbf{k}(t)\rangle. \quad (10)$$

Here k_i denotes the initial momentum at $t = 0$. That is, $k_1(0) = 0$, $k_2(0) = 0$ and $k_3(0) = k \cos\theta(0) = k$. Thus it follows from Eq.(10) that

$$\mathbf{p}|m, \mathbf{k}(t)\rangle = \mathbf{k}(t)|m, \mathbf{k}(t)\rangle, \quad (11)$$

which is shown to be the eigenvalue equation of the momentum operator of atoms in the curved atom fiber. Thus, the solution $|m, \mathbf{k}(t)\rangle$ of Schrödinger equation (5) is truly the eigenstate of the momentum operator $\mathbf{p}$ of atoms.

In addition, the operator $I(t) = \mathbf{k}(t) \cdot \mathbf{J}/k$, which is the total angular momentum operator, $\mathbf{J}$, of atoms projected onto the atom momentum (wave vector) $\mathbf{k}(t)$, is a physically interesting quantity, since it agrees with the Liouville-Von Neumann equation $\partial I(t)/\partial t - i[I(t), H(t)] = 0$. This, therefore, means that the operator $\mathbf{k}(t) \cdot \mathbf{J}/k$ can be considered a Lewis-Riesenfeld invariant [13], which possesses the conserved eigenvalues in the evolution process. Thus, for the atomic matter waves in the atom fiber, it may be referred to as the atomic *helicity operator*.

To conclude, assuming that the atomic matter wave is propagating inside a perfect atom guide, we construct an effective Hamiltonian to describe the interaction between the atomic matter waves and the spiral shaped guide, and present the exact solutions of the time-dependent Schrödinger equation that governs the evolution of matter wave in the coiled atom fiber (guide), as realized experimentally by Luo *et al.* [7]. Since it is shown that such wavefunctions is truly the eigenstates of momentum operator of atoms, we think that, at least within the framework of the two assumptions made above, our method is self-consistent. So, it may be believed that the formulation presented here would provide us with a useful insight into the evolution problem in the atom fibers, and therefore deserves further consideration in the investigation of atom optics.

Acknowledgements This project was supported partially by the National Natural Science Foundation of China under Project No. 90101024 and 60378037. I thank Q. Li and D. Wang for their valuable suggestions and comments on my manuscript.

-
- [1] R. Folman, P. Krüger, J. Schmiedmayer, J. Denschlag, and C. Henkel, *Adv. At. Mol. Opt. Phys.* **48**, 263 (2002).
 - [2] J. Reichel, *Appl. Phys. B* **74**, 469 (2002).
 - [3] P. Krüger, X. Luo, M.W. Klein *et al.*, arXiv: quant-ph/0306111 (2003).
 - [4] J. Reichel, W. Hänsel, and T.W. Hänsch, *Phys. Rev. Lett.* **83**, 3398 (1999).
 - [5] R. Folman, P. Krüger, D. Cassettari, B. Hessmo, T. Maier, and J. Schmiedmayer, *Phys. Rev. Lett.* **84**, 4749 (2000).
 - [6] D. Cassettari, B. Hessmo, R. Folman, T. Maier, and J. Schmiedmayer, *Phys. Rev. Lett.* **85**, 5483 (2000).
 - [7] X. Luo, P. Krüger, K. Brugger *et al.*, arXiv: quant-ph/0311174 (2003).
 - [8] J.Q. Shen, H.Y. Zhu, S.L. Shi, and J. Li, *Phys. Scr.* **65**, 465 (2002).
 - [9] J.Q. Shen and S.L. He, *Phys. Rev. B* **68**, 195421 (2003).
 - [10] B. Mashhoon, *Phys. Lett. A* **173**, 347 (1993).
 - [11] B. Mashhoon, *Gen. Relativ. Gravit.* **31**, 681 (1999).
 - [12] J.Q. Shen, H.Y. Zhu, and P. Chen, *Eur. Phys. J. D* **23**, 305 (2003).
 - [13] H.R. Lewis and W.B. Riesenfeld, *J. Math. Phys.* **10**, 1458 (1969).